

preprint: May 30, 2016

New super congruences involving Bernoulli and Euler polynomials

Zhi-Hong Sun

School of Mathematical Sciences
Huaiyin Normal University
Huaian, Jiangsu 223001, P.R. China
zhsun@hytc.edu.cn
<http://www.hytc.edu.cn/xsjl/szh>

Let $p > 3$ be a prime, and let a be a rational p -adic integer with $a \not\equiv 0 \pmod{p}$. In this paper we establish congruences for

$$\sum_{k=1}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{k}, \quad \sum_{k=0}^{(p-1)/2} k \binom{a}{k} \binom{-1-a}{k} \quad \text{and} \quad \sum_{k=0}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{2k-1} \pmod{p^2}$$

in terms of Bernoulli and Euler polynomials. We also give some transformation formulas for congruences modulo p^2 .

Keywords: Congruence; Bernoulli polynomial; Euler polynomial.

Mathematics Subject Classification 2010: Primary 11A07, Secondary 11B68, 05A10, 05A19, 11B39

1. Introduction

The Bernoulli numbers $\{B_n\}$ and Bernoulli polynomials $\{B_n(x)\}$ are defined by

$$B_0 = 1, \quad \sum_{k=0}^{n-1} \binom{n}{k} B_k = 0 \quad (n \geq 2) \quad \text{and} \quad B_n(x) = \sum_{k=0}^n \binom{n}{k} B_k x^{n-k} \quad (n \geq 0).$$

The Euler numbers $\{E_n\}$ and Euler polynomials $\{E_n(x)\}$ are defined by

$$E_0 = 1, \quad E_n = - \sum_{k=1}^{[n/2]} \binom{n}{2k} E_{n-2k} \quad (n \geq 1) \quad \text{and} \quad E_n(x) = \frac{1}{2^n} \sum_{r=0}^n \binom{n}{r} (2x-1)^{n-r} E_r \quad (n \geq 0),$$

where $[a]$ is the greatest integer not exceeding a . In [S6] the author introduced the sequence $\{U_n\}$ given by

$$U_0 = 1 \quad \text{and} \quad U_n = -2 \sum_{k=1}^{[n/2]} \binom{n}{2k} U_{n-2k} \quad (n \geq 1).$$

It is well known that $B_{2n+1} = 0$ and $E_{2n-1} = U_{2n-1} = 0$ for any positive integer n . $\{B_n\}$, $\{E_n\}$ and $\{U_n\}$ are important sequences and they have many interesting properties and applications. See [EMOT], [IR], [MOS] and [S2-S7].

It is easily seen that ([S8])

$$(1.1) \quad \begin{aligned} \binom{-\frac{1}{2}}{k}^2 &= \frac{\binom{2k}{k}^2}{16^k}, \quad \binom{-\frac{1}{3}}{k} \binom{-\frac{2}{3}}{k} = \frac{\binom{2k}{k} \binom{3k}{k}}{27^k}, \\ \binom{-\frac{1}{4}}{k} \binom{-\frac{3}{4}}{k} &= \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k}, \quad \binom{-\frac{1}{6}}{k} \binom{-\frac{5}{6}}{k} = \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k}. \end{aligned}$$

In 2003, Rodriguez-Villegas [RV] conjectured that for any prime $p > 3$,

$$(1.2) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k} \equiv \left(\frac{-1}{p}\right) \pmod{p^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} \equiv \left(\frac{-3}{p}\right) \pmod{p^2},$$

$$(1.3) \quad \sum_{k=0}^{p-1} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k} \equiv \left(\frac{-2}{p}\right) \pmod{p^2}, \quad \sum_{k=0}^{p-1} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k} \equiv \left(\frac{-1}{p}\right) \pmod{p^2},$$

where $\left(\frac{a}{p}\right)$ is the Legendre symbol. (1.2) and (1.3) were later confirmed by Mortenson [M1-M2]. Let $\mathbb{Z}$ be the set of integers. For a prime p let $\mathbb{Z}_p$ denote the set of rational numbers whose denominator is not divisible by p . For $a \in \mathbb{Z}_p$ let $\langle a \rangle_p \in \{0, 1, \dots, p-1\}$ be given by $a \equiv \langle a \rangle_p \pmod{p}$. Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$, $a \not\equiv 0 \pmod{p}$ and $t = (a - \langle a \rangle_p)/p$. In [S9-S11] the author showed that

$$(1.4) \quad \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \equiv (-1)^{\langle a \rangle_p} + p^2 t(t+1) E_{p-3}(-a) \pmod{p^3},$$

$$(1.5) \quad \sum_{k=0}^{p-1} k \binom{a}{k} \binom{-1-a}{k} \equiv (-1)^{\langle a \rangle_p} a(a+1) + p^2 t(t+1) (a(a+1) E_{p-3}(-a) - 1) \pmod{p^3},$$

$$(1.6) \quad \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{1}{2k-1} \equiv -(2a+1)(2t+1) - p^2 t(t+1)(4 + (2a+1) B_{p-2}(-a)) \pmod{p^3},$$

$$(1.7) \quad \sum_{k=0}^{p-1} \binom{a}{k} \binom{-1-a}{k} \frac{2a+1}{2k+1} \equiv 1 + 2t + p^2 t(t+1) B_{p-2}(-a) \pmod{p^3},$$

$$(1.8) \quad \sum_{k=1}^{p-1} \frac{1}{k} \binom{a}{k} \binom{-1-a}{k} \equiv -\frac{2}{3} p^2 t(t+1) B_{p-3}(-a) - 2 \frac{B_{p^2(p-1)}(-a) - B_{p^2(p-1)}}{p^2(p-1)} \pmod{p^3},$$

$$(1.9) \quad \sum_{k=0}^{p-1} \binom{a}{k} (-2)^k \equiv (-1)^{\langle a \rangle_p} - p t E_{p-2}(-a) \pmod{p^2},$$

$$(1.10) \quad \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{a}{k} \equiv \frac{B_{p^2(p-1)}(-a) - B_{p^2(p-1)}}{p^2(p-1)} - \frac{a - \langle a \rangle_p}{2} B_{p-2}(-a) \pmod{p^2}.$$

We note that Tauraso[T2] obtained a congruence for $\sum_{k=1}^{p-1} \frac{1}{k} \binom{a}{k} \binom{-1-a}{k} \pmod{p^2}$, and he proved that ([T1])

$$\sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{-1/2}{k} \equiv H_{\frac{p-1}{2}} \pmod{p^3},$$

where $H_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n}$. Z.W. Sun proved that ([Su1])

$$\sum_{k=0}^{p-1} \frac{\binom{2k}{k}^2}{16^k} = \sum_{k=0}^{p-1} \binom{-\frac{1}{2}}{k}^2 \equiv 1 + (-1)^{\frac{p-1}{2}} - p^2 E_{p-3} \pmod{p^3}.$$

Recently, using the fact

$$(1.11) \quad \binom{a-1}{\frac{p-1}{2}} \binom{-a-1}{\frac{p-1}{2}} \equiv \begin{cases} \frac{pt}{\langle a \rangle_p} \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ \frac{p(t+1)}{\langle a \rangle_p} \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2} \end{cases}$$

and the method in [S9-S11] Mao and Sun [MS] obtained congruences for $\sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k}$ and $\sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} \frac{1}{2k+1} \pmod{p^2}$. In particular, they proved

$$\begin{aligned} \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{3k}{k}}{27^k} &\equiv \left(\frac{p}{3}\right) \frac{2^p + 1}{3} \pmod{p^2}, \quad \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{4k}{2k}}{(2k+1)64^k} \equiv \left(\frac{-1}{p}\right) 2^{p-1} \pmod{p^2}, \\ \sum_{k=0}^{(p-1)/2} \frac{\binom{3k}{k} \binom{6k}{3k}}{(2k+1)432^k} &\equiv \left(\frac{p}{3}\right) \frac{3^p + 1}{4} \pmod{p^2}. \end{aligned}$$

Let $p > 3$ be a prime and $a \in \mathbb{Z}_p$ with $a \not\equiv 0 \pmod{p}$. In this paper we establish congruences for

$$\sum_{k=1}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{k}, \quad \sum_{k=0}^{(p-1)/2} k \binom{a}{k} \binom{-1-a}{k} \quad \text{and} \quad \sum_{k=0}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{2k-1} \pmod{p^2}$$

and some transformation formulas for congruences modulo p^2 . For instance, for $\langle a \rangle_p < \frac{p}{2}$ we have

$$\begin{aligned} \sum_{k=1}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{k} &\equiv -2 \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{a}{k} \pmod{p^2} \\ \sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} &\equiv \sum_{k=0}^{p-1} \binom{a}{k} (-2)^k \pmod{p^2}, \\ \sum_{k=0}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{2k-1} &\equiv 2(a - \langle a \rangle_p) - (2a+1) \sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \pmod{p^2}. \end{aligned}$$

As consequences and applications we obtain some new congruences modulo p^2 . Here are three typical examples:

$$(1.12) \quad \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{4k}{2k}}{(2k-1)64^k} \equiv (-1)^{\frac{p+1}{2}} \frac{p+1}{2} \pmod{p^2},$$

$$(1.13) \quad \sum_{k=1}^{(p-1)/2} \frac{\binom{2k}{k} \binom{4k}{2k}}{k \cdot 64^k} \equiv 6q_p(2) - p(3q_p(2)^2 + 2(-1)^{\frac{p-1}{2}} E_{p-3}) \pmod{p^2},$$

$$(1.14) \quad \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k} \equiv (-1)^{[\frac{p}{4}]} (1 + P_{p-(\frac{2}{p})}) \pmod{p^2}.$$

where $q_p(a) = (a^{p-1} - 1)/p$ and $\{P_n\}$ is the Pell sequence given by $P_0 = 0$, $P_1 = 1$ and $P_{n+1} = 2P_n + P_{n-1}$ ($n \geq 1$). We note that (1.13) is equivalent to a conjecture made by the author's brother Z.W. Sun in [Su2].

2. Congruences for $\sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k} \binom{a}{k} \binom{-1-a}{k} \pmod{p^2}$

For any positive integer n and variable a let

$$S_n(a) = \sum_{k=1}^n \frac{1}{k} \binom{a}{k} \binom{-1-a}{k}.$$

By [S9, (2.1)],

$$(2.1) \quad S_n(a) - S_n(a-1) = \frac{2}{a} \binom{a-1}{n} \binom{-a-1}{n} - \frac{2}{a}.$$

Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$ and $t = (a - \langle a \rangle_p)/p$. Using (2.1) we see that

$$\begin{aligned} & S_n(a) - S_n(a - \langle a \rangle_p) \\ &= \sum_{k=0}^{\langle a \rangle_p - 1} (S_n(a-k) - S_n(a-k-1)) = \sum_{k=0}^{\langle a \rangle_p - 1} \frac{2}{a-k} \left\{ \binom{a-k-1}{n} \binom{k-a-1}{n} - 1 \right\} \\ &= \sum_{k=0}^{\langle a \rangle_p - 1} \frac{2}{pt + \langle a \rangle_p - k} \left\{ \binom{pt + \langle a \rangle_p - k - 1}{n} \binom{-pt - (\langle a \rangle_p - k) - 1}{n} - 1 \right\}. \end{aligned}$$

Hence

$$(2.2) \quad S_n(a) - S_n(pt) = \sum_{r=1}^{\langle a \rangle_p} \frac{2}{pt+r} \left\{ \binom{pt+r-1}{n} \binom{-pt-r-1}{n} - 1 \right\}.$$

For any positive integer n define $H_n = \sum_{k=1}^n \frac{1}{k}$. For convenience we also define $H_0 = 0$.

Lemma 2.1. *Let $p > 3$ be a prime, $r \in \{1, 2, \dots, p-1\}$ and $t \in \mathbb{Z}_p$. Then*

$$\begin{aligned} & \binom{pt+r-1}{\frac{p-1}{2}} \binom{-pt-r-1}{\frac{p-1}{2}} \\ & \equiv \begin{cases} \frac{pt}{r} + \frac{p^2 t}{r} (2q_p(2) + H_{\frac{p-1}{2}-r}) - \frac{p^2 t^2}{r^2} \pmod{p^3} & \text{if } r < \frac{p}{2}, \\ \frac{p(t+1)}{r} + \frac{p^2(t+1)}{r} (2q_p(2) + H_{r-\frac{p+1}{2}}) - \frac{p^2 t(t+1)}{r^2} \pmod{p^3} & \text{if } r > \frac{p}{2}. \end{cases} \end{aligned}$$

Proof. We first assume $r < \frac{p}{2}$. It is clear that

$$\begin{aligned} & \left(\frac{p-1}{2}\right)! \binom{pt+r-1}{\frac{p-1}{2}} \binom{-pt-r-1}{\frac{p-1}{2}} \\ & = \left(\frac{p-1}{2}\right)! \binom{pt+r-1}{\frac{p-1}{2}} (-1)^{\frac{p-1}{2}} \binom{pt+r+\frac{p-1}{2}}{\frac{p-1}{2}} \\ & = (-1)^{\frac{p-1}{2}} \frac{pt}{pt+r} (pt-1) \cdots \left(pt - \left(\frac{p-1}{2} - r\right)\right) (pt+1)(pt+2) \cdots \left(pt + \frac{p-1}{2} + r\right) \\ & = (-1)^{\frac{p-1}{2}} \frac{pt}{pt+r} ((pt)^2 - 1^2)((pt)^2 - 2^2) \cdots \left((pt)^2 - \left(\frac{p-1}{2} - r\right)^2\right) \\ & \quad \times \prod_{s=0}^{r-1} \left(pt + \frac{p-1}{2} - s\right) \left(pt + \frac{p-1}{2} + s + 1\right) \\ & \equiv (-1)^r \frac{pt}{pt+r} \left(\frac{p-1}{2} - r\right)! \prod_{s=0}^{r-1} \left(\frac{p-1}{2} - s\right) \left(\frac{p-1}{2} + s + 1\right) \\ & = (-1)^r \frac{pt}{pt+r} \left(\frac{p-1}{2} - r\right)! \left(\frac{p-1}{2} + r\right)! = (-1)^r \frac{pt}{pt+r} \cdot \frac{(p-1)!}{\left(\frac{p-1}{2}-r\right)!} \\ & \equiv (-1)^r \frac{pt}{pt+r} \cdot \frac{(p-1)!}{(-1)^{\frac{p-1}{2}-r} (1 - pH_{\frac{p-1}{2}-r})} \equiv (-1)^{\frac{p-1}{2}} (p-1)! \cdot pt \frac{(r-pt)(1 + pH_{\frac{p-1}{2}-r})}{r^2} \\ & \equiv (-1)^{\frac{p-1}{2}} (p-1)! \cdot pt \left(\frac{1}{r} - \frac{pt}{r^2} + \frac{pH_{\frac{p-1}{2}-r}}{r}\right) \pmod{p^3}. \end{aligned}$$

Since

$$\frac{(p-1)!}{\left(\frac{p-1}{2}\right)!^2} = \frac{(p-1) \cdots (p - \frac{p-1}{2})}{\left(\frac{p-1}{2}\right)!} \equiv (-1)^{\frac{p-1}{2}} (1 - pH_{\frac{p-1}{2}}) \equiv (-1)^{\frac{p-1}{2}} (1 + 2pq_p(2)) \pmod{p^2},$$

from the above we deduce that

$$\begin{aligned} \binom{pt+r-1}{\frac{p-1}{2}} \binom{-pt-r-1}{\frac{p-1}{2}} & \equiv (1 + 2pq_p(2)) pt \left(\frac{1}{r} - \frac{pt}{r^2} + \frac{pH_{\frac{p-1}{2}-r}}{r}\right) \\ & \equiv pt \left(\frac{1}{r} + \frac{p(2q_p(2) + H_{\frac{p-1}{2}-r})}{r} - \frac{pt}{r^2}\right) \pmod{p^3}. \end{aligned}$$

This yields the result in the case $r < \frac{p}{2}$.

Now assume $r > \frac{p}{2}$. Set $t' = -t - 1$ and $r' = p - r$. We see that

$$\begin{aligned} \binom{pt+r-1}{\frac{p-1}{2}} \binom{-pt-r-1}{\frac{p-1}{2}} &= \binom{p(t+1)-(p-r)-1}{\frac{p-1}{2}} \binom{-p(t+1)+p-r-1}{\frac{p-1}{2}} \\ &= \binom{pt'+r'-1}{\frac{p-1}{2}} \binom{-pt'-r'-1}{\frac{p-1}{2}}. \end{aligned}$$

Since $r' < \frac{p}{2}$, from the above we deduce that

$$\begin{aligned} &\binom{pt+r-1}{\frac{p-1}{2}} \binom{-pt-r-1}{\frac{p-1}{2}} \\ &= \binom{pt'+r'-1}{\frac{p-1}{2}} \binom{-pt'-r'-1}{\frac{p-1}{2}} \equiv \frac{pt'}{r'} + \frac{p^2 t'}{r'} (2q_p(2) + H_{\frac{p-1}{2}-r'}) - \frac{p^2 t'^2}{r'^2} \\ &= \frac{p(t+1)}{r-p} - \frac{p^2(t+1)}{p-r} (2q_p(2) + H_{r-\frac{p+1}{2}}) - \frac{p^2(t+1)^2}{(p-r)^2} \\ &\equiv \frac{p(t+1)(r+p)}{r^2} + \frac{p^2(t+1)}{r} (2q_p(2) + H_{r-\frac{p+1}{2}}) - \frac{p^2(t+1)^2}{r^2} \pmod{p^3}. \end{aligned}$$

This yields the result in the case $r > \frac{p}{2}$. Hence the lemma is proved.

We remark that Lemma 2.1 improves Mao and Sun's (1.11).

Theorem 2.1. *Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$ and $a \not\equiv 0 \pmod{p}$. Then*

$$\sum_{k=1}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{k} + 2 \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{a}{k} \equiv \begin{cases} 0 \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ pB_{p-2}(-a) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2} \end{cases}$$

and

$$\begin{aligned} &\sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{a}{k} \\ &\equiv -\frac{B_{2p-2}(-a) - B_{2p-2}}{2p-2} + 2\frac{B_{p-1}(-a) - B_{p-1}}{p-1} - \frac{a - \langle a \rangle_p}{2} B_{p-2}(-a) \pmod{p^2}. \end{aligned}$$

Proof. Set $t = (a - \langle a \rangle_p)/p$. By [S3, Theorem 5.2], $\sum_{k=1}^{(p-1)/2} \frac{1}{k^2} \equiv 0 \pmod{p}$. Thus,

$$\begin{aligned} S_{\frac{p-1}{2}}(pt) &= \sum_{k=1}^{(p-1)/2} \frac{1}{k} \binom{pt}{k} \binom{-pt-1}{k} = \sum_{k=1}^{(p-1)/2} \frac{pt}{k^2} \binom{pt-1}{k-1} \binom{-pt-1}{k} \\ &\equiv \sum_{k=1}^{(p-1)/2} \frac{pt}{k^2} \binom{-1}{k-1} \binom{-1}{k} = -pt \sum_{k=1}^{(p-1)/2} \frac{1}{k^2} \equiv 0 \pmod{p^2}. \end{aligned}$$

For $1 \leq \langle a \rangle_p \leq \frac{p-1}{2}$, from (2.2) and Lemma 2.1 we see that

$$S_{\frac{p-1}{2}}(a) = S_{\frac{p-1}{2}}(pt) + \sum_{r=1}^{\langle a \rangle_p} \frac{2(r-pt)}{r^2 - p^2 t^2} \left\{ \binom{pt+r-1}{\frac{p-1}{2}} \binom{-pt-r-1}{\frac{p-1}{2}} - 1 \right\}$$

$$\equiv 2 \sum_{r=1}^{\langle a \rangle_p} \frac{r-pt}{r^2} \left(\frac{pt}{r} - 1 \right) \equiv 2 \sum_{r=1}^{\langle a \rangle_p} \left(\frac{2pt}{r^2} - \frac{1}{r} \right) \pmod{p^2}.$$

For $\langle a \rangle_p > \frac{p-1}{2}$, from (2.2) and Lemma 2.1 we deduce that

$$\begin{aligned} S_{\frac{p-1}{2}}(a) &= S_{\frac{p-1}{2}}(pt) + \sum_{r=1}^{\langle a \rangle_p} \frac{2(r-pt)}{r^2 - p^2 t^2} \left\{ \binom{pt+r-1}{\frac{p-1}{2}} \binom{-pt-r-1}{\frac{p-1}{2}} - 1 \right\} \\ &\equiv 2 \sum_{r=1}^{(p-1)/2} \frac{r-pt}{r^2} \left(\frac{pt}{r} - 1 \right) + 2 \sum_{r=(p+1)/2}^{\langle a \rangle_p} \frac{r-pt}{r^2} \left(\frac{p(t+1)}{r} - 1 \right) \\ &= 2 \sum_{r=1}^{\langle a \rangle_p} \frac{r-pt}{r^2} \left(\frac{p(t+1)}{r} - 1 \right) - 2 \sum_{r=1}^{(p-1)/2} \frac{r-pt}{r^2} \cdot \frac{p}{r} \\ &\equiv 2 \sum_{r=1}^{\langle a \rangle_p} \left(\frac{2pt}{r^2} - \frac{1}{r} \right) + 2p \sum_{r=1}^{\langle a \rangle_p} \frac{1}{r^2} - 2p \sum_{r=1}^{(p-1)/2} \frac{1}{r^2} \pmod{p^2}. \end{aligned}$$

By [S3, Theorem 5.2], $\sum_{r=1}^{(p-1)/2} \frac{1}{r^2} \equiv 0 \pmod{p}$. By [S9, (3.4)] and Fermat's little theorem, $\sum_{r=1}^{\langle a \rangle_p} \frac{1}{r^2} \equiv \frac{1}{2} B_{p-2}(-a) \pmod{p}$. From the above we deduce that

$$(2.3) \quad S_{\frac{p-1}{2}}(a) - 2 \sum_{r=1}^{\langle a \rangle_p} \left(\frac{2pt}{r^2} - \frac{1}{r} \right) \equiv \begin{cases} 0 \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ p B_{p-2}(-a) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2}. \end{cases}$$

Putting $n = p-1$ and $b = -1$ in [S9, Lemma 3.1] and then applying [S3, Theorem 5.1] we get

$$\sum_{k=1}^{p-1} \frac{(-1)^k}{k} \binom{a}{k} \equiv -pt \sum_{k=1}^{p-1} \frac{1}{k^2} - \sum_{r=1}^{\langle a \rangle_p} \frac{1}{r} + 2pt \sum_{r=1}^{\langle a \rangle_p} \frac{1}{r^2} \equiv \sum_{r=1}^{\langle a \rangle_p} \left(\frac{2pt}{r^2} - \frac{1}{r} \right) \pmod{p^2}.$$

This together with (2.3) yields the the first part.

Putting $b = p-1$ and $x = -a$ in [S3, Theorem 3.1] and then applying [S3, Theorem 2.2 (with $k = p^2 - 1$ and $n = 2$)] we deduce that

$$\begin{aligned} \frac{B_{p^2(p-1)}(-a) - B_{p^2(p-1)}}{p^2(p-1)} &= \frac{B_{(p^2-1)(p-1)+p-1}(-a) - B_{(p^2-1)(p-1)+p-1}}{(p^2-1)(p-1) + p-1} \\ &\equiv (p^2-1) \frac{B_{p-1+p-1}(-a) - B_{p-1+p-1}}{p-1+p-1} - (p^2-2) \frac{B_{p-1}(-a) - B_{p-1}}{p-1} \\ &\equiv -\frac{B_{2p-2}(-a) - B_{2p-2}}{2p-2} + 2 \frac{B_{p-1}(-a) - B_{p-1}}{p-1} \pmod{p^2}. \end{aligned}$$

This together with (1.10) ([S9, Theorem 3.1]) yields the remaining part. The proof is now complete.

Theorem 2.2. *Let $p > 3$ be a prime. Then*

- (i)
$$\sum_{k=1}^{(p-1)/2} \frac{\binom{2k}{k} \binom{4k}{2k}}{k \cdot 64^k} \equiv 6q_p(2) - p(3q_p(2)^2 + 2(-1)^{\frac{p-1}{2}} E_{p-3}) \pmod{p^2},$$
- (ii)
$$\sum_{k=1}^{(p-1)/2} \frac{\binom{2k}{k} \binom{3k}{k}}{k \cdot 27^k} \equiv 3q_p(3) - p\left(\frac{3}{2}q_p(3)^2 + 2\left(\frac{p}{3}\right)U_{p-3}\right) \pmod{p^2},$$
- (iii)
$$\sum_{k=1}^{(p-1)/2} \frac{\binom{3k}{k} \binom{6k}{3k}}{k \cdot 432^k} \equiv 4q_p(2) + 3q_p(3) - p\left(2q_p(2)^2 + \frac{3}{2}q_p(3)^2 + 5\left(\frac{p}{3}\right)U_{p-3}\right) \pmod{p^2}.$$

Proof. From [S4, Lemma 2.5] we know that $E_{2n} = -4^{2n+1} \frac{B_{2n+1}(\frac{1}{4})}{2n+1}$. Thus, $E_{p-3} = -4^{p-2} \frac{B_{p-2}(\frac{1}{4})}{p-2} \equiv \frac{1}{8}B_{p-2}(\frac{1}{4}) \pmod{p}$. Now taking $a = -\frac{1}{4}$ in Theorem 2.1 we see that

$$\begin{aligned} & \sum_{k=1}^{(p-1)/2} \frac{\binom{-1/4}{k} \binom{-3/4}{k}}{k} + 2 \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{-1/4}{k} \\ & \equiv \begin{cases} 0 \pmod{p} & \text{if } p \equiv 1 \pmod{4}, \\ pB_{p-2}(\frac{1}{4}) \equiv 8pE_{p-3} \pmod{p} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

By [S9, Theorem 3.2],

$$-2 \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{-1/4}{k} \equiv 6q_p(2) + p(-3q_p(2)^2 - 2(2 - (-1)^{\frac{p-1}{2}})E_{p-3}) \pmod{p^2}.$$

Hence (i) is true by (1.1).

Taking $a = -\frac{1}{3}$ in Theorem 2.1 and applying the fact $B_{p-2}(\frac{1}{3}) \equiv 6U_{p-3} \pmod{p}$ ([S6]) we see that

$$\begin{aligned} & \sum_{k=1}^{(p-1)/2} \frac{\binom{-1/3}{k} \binom{-2/3}{k}}{k} + 2 \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{-1/3}{k} \\ & \equiv \begin{cases} 0 \pmod{p} & \text{if } p \equiv 1 \pmod{3}, \\ pB_{p-2}(\frac{1}{3}) \equiv 6pU_{p-3} \pmod{p} & \text{if } p \equiv 2 \pmod{3}. \end{cases} \end{aligned}$$

By [S9, Theorem 3.3],

$$-2 \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{-1/3}{k} \equiv 3q_p(3) - p\left(\frac{3}{2}q_p(3)^2 + (3 - \frac{p}{3})U_{p-3}\right) \pmod{p^2}.$$

Thus (ii) holds by (1.1).

Taking $a = -\frac{1}{6}$ in Theorem 2.1 and applying the fact $B_{p-2}(\frac{1}{6}) \equiv 30U_{p-3} \pmod{p}$ ([S6]) we see that

$$\sum_{k=1}^{(p-1)/2} \frac{\binom{-1/6}{k} \binom{-5/6}{k}}{k} + 2 \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{-1/6}{k}$$

$$\equiv \begin{cases} 0 \pmod{p} & \text{if } p \equiv 1 \pmod{3}, \\ pB_{p-2}\left(\frac{1}{6}\right) \equiv 30pU_{p-3} \pmod{p} & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

By [S9, Theorem 3.4],

$$\begin{aligned} & -2 \sum_{k=1}^{p-1} \frac{(-1)^{k-1}}{k} \binom{-1/6}{k} \\ & \equiv 4q_p(2) + 3q_p(3) - p \left(2q_p(2)^2 + \frac{3}{2}q_p(3)^2 + 5\left(3 - 2\left(\frac{p}{3}\right)\right)U_{p-3} \right) \pmod{p^2}. \end{aligned}$$

Thus (iii) holds by (1.1).

3. Congruences for $\sum_{k=0}^{\frac{p-1}{2}} \binom{a}{k} \binom{-1-a}{k} \pmod{p^2}$ and $\sum_{k=0}^{\frac{p-1}{2}} k \binom{a}{k} \binom{-1-a}{k} \pmod{p^2}$

Let p be a prime greater than 3, $a \in \mathbb{Z}_p$ and $t = (a - \langle a \rangle_p)/p$. From [S11, p.3299] we know that for any positive integer n ,

$$\sum_{k=0}^n \binom{a}{k} \binom{-1-a}{k} - (-1)^{\langle a \rangle_p} \sum_{k=0}^n \binom{pt}{k} \binom{-1-pt}{k} = 2 \sum_{k=0}^{\langle a \rangle_p - 1} (-1)^k \binom{a-k-1}{n} \binom{k-a-1}{n}.$$

For $n < p$ we see that

$$\begin{aligned} \sum_{k=0}^n \binom{pt}{k} \binom{-1-pt}{k} &= 1 + \sum_{k=1}^n \frac{pt}{k} \binom{pt-1}{k-1} \binom{-1-pt}{k} \\ &\equiv 1 + pt \sum_{k=1}^n \frac{1}{k} \binom{-1}{k-1} \binom{-1}{k} = 1 - ptH_n \pmod{p^2}. \end{aligned}$$

Taking $n = \frac{p-1}{2}$ in the above and then applying (1.11) or Lemma 2.1 we deduce the following lemma due to Mao and Sun.

Lemma 3.1 ([MS, (3.5)]. *Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$, $a \not\equiv 0 \pmod{p}$ and $t = (a - \langle a \rangle_p)/p$. Then*

$$\sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} \equiv (-1)^{\langle a \rangle_p} (1 - ptH_{\frac{p-1}{2}}) + 2p \sum_{k=0}^{\langle a \rangle_p - 1} (-1)^k \frac{t + \delta_k}{\langle a \rangle_p - k} \pmod{p^2},$$

where $\delta_k = 1$ or 0 according as $\langle a \rangle_p - k > \frac{p-1}{2}$ or not.

Theorem 3.1. *Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$, $a \not\equiv 0 \pmod{p}$ and $t = (a - \langle a \rangle_p)/p$. Then*

$$\sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k}$$

$$\equiv \begin{cases} \sum_{k=0}^{p-1} \binom{a}{k} (-2)^k \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ \left(1 + \frac{1}{t}\right) \sum_{k=0}^{p-1} \binom{a}{k} (-2)^k - (-1)^{\langle a \rangle_p} \frac{1}{t} \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2} \text{ and } t \not\equiv 0 \pmod{p}, \\ (-1)^{\langle a \rangle_p} - pE_{p-2}(-a) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2} \text{ and } t \equiv 0 \pmod{p}. \end{cases}$$

Proof. Note that

$$\begin{aligned} \sum_{k=0}^{\langle a \rangle_p - 1} (-1)^{\langle a \rangle_p - k} \frac{t + \delta_k}{\langle a \rangle_p - k} &= \sum_{\substack{0 \leq k \leq \langle a \rangle_p - 1 \\ \langle a \rangle_p - k < \frac{p}{2}}} (-1)^{\langle a \rangle_p - k} \frac{t}{\langle a \rangle_p - k} + \sum_{\substack{0 \leq k \leq \langle a \rangle_p - 1 \\ \langle a \rangle_p - k > \frac{p}{2}}} (-1)^{\langle a \rangle_p - k} \frac{t+1}{\langle a \rangle_p - k} \\ &= \sum_{\substack{1 \leq r \leq \langle a \rangle_p \\ r < \frac{p}{2}}} (-1)^r \frac{t}{r} + \sum_{\substack{1 \leq r \leq \langle a \rangle_p \\ r > \frac{p}{2}}} (-1)^r \frac{t+1}{r} \\ &= \begin{cases} \sum_{r=1}^{\langle a \rangle_p} (-1)^r \frac{t}{r} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ \sum_{r=1}^{\langle a \rangle_p} (-1)^r \frac{t+1}{r} - \sum_{r=1}^{(p-1)/2} \frac{(-1)^r}{r} & \text{if } \langle a \rangle_p > \frac{p-1}{2}. \end{cases} \end{aligned}$$

From [L] we know that $H_{\frac{p-1}{2}} \equiv -2q_p(2) \pmod{p}$ and $H_{[\frac{p}{4}]} \equiv -3q_p(2) \pmod{p}$. Thus,

$$\sum_{r=1}^{(p-1)/2} \frac{(-1)^r}{r} = \sum_{r=1}^{(p-1)/2} \frac{(-1)^r + 1}{r} - H_{\frac{p-1}{2}} = H_{[\frac{p}{4}]} - H_{\frac{p-1}{2}} \equiv -q_p(2) \pmod{p}.$$

Hence applying Lemma 3.1 we obtain

$$\begin{aligned} \sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} &\equiv (-1)^{\langle a \rangle_p} (1 + 2ptq_p(2)) + 2pt(-1)^{\langle a \rangle_p} \sum_{r=1}^{\langle a \rangle_p} \frac{(-1)^r}{r} \\ &+ \begin{cases} 0 \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ 2p(-1)^{\langle a \rangle_p} \sum_{r=1}^{\langle a \rangle_p} \frac{(-1)^r}{r} + 2p(-1)^{\langle a \rangle_p} q_p(2) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2}. \end{cases} \end{aligned}$$

By [S11, p.3306],

$$\sum_{r=1}^{\langle a \rangle_p} \frac{(-1)^r}{r} + q_p(2) \equiv -\frac{1}{2}(-1)^{\langle a \rangle_p} E_{p-2}(-a) \pmod{p}.$$

Thus,
(3.1)

$$\sum_{k=0}^{\frac{p-1}{2}} \binom{a}{k} \binom{-1-a}{k} \equiv \begin{cases} (-1)^{\langle a \rangle_p} (1 + 2pt \left(\sum_{r=1}^{\langle a \rangle_p} \frac{(-1)^r}{r} + q_p(2) \right)) \\ \equiv (-1)^{\langle a \rangle_p} - ptE_{p-2}(-a) \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ (-1)^{\langle a \rangle_p} (1 + 2p(t+1) \left(\sum_{r=1}^{\langle a \rangle_p} \frac{(-1)^r}{r} + q_p(2) \right)) \\ \equiv (-1)^{\langle a \rangle_p} - p(t+1)E_{p-2}(-a) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2}. \end{cases}$$

By [S11, Theorem 3.1],

$$(-1)^{\langle a \rangle_p} - ptE_{p-2}(-a) \equiv \sum_{k=0}^{p-1} \binom{a}{k} (-2)^k \pmod{p^2}.$$

Thus, the result is true for $\langle a \rangle_p \leq \frac{p-1}{2}$. For $\langle a \rangle_p > \frac{p-1}{2}$ and $t \not\equiv 0 \pmod{p}$ we have

$$\begin{aligned} \sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} &\equiv (-1)^{\langle a \rangle_p} + \frac{t+1}{t} (-ptE_{p-2}(-a)) \\ &\equiv (-1)^{\langle a \rangle_p} + \frac{t+1}{t} \left(\sum_{k=0}^{p-1} \binom{a}{k} (-2)^k - (-1)^{\langle a \rangle_p} \right) \pmod{p^2}. \end{aligned}$$

This yields the result in this case. The proof is now complete.

Theorem 3.2. *Let $p > 3$ be a prime. Then*

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k} \equiv (-1)^{\lfloor \frac{p}{4} \rfloor} (1 + P_{p-(\frac{2}{p})}) \pmod{p^2}.$$

Proof. Clearly

$$\sum_{r=1}^{\langle -1/4 \rangle_p} \frac{(-1)^r}{r} = \sum_{r=1}^{\langle -1/4 \rangle_p} \frac{(-1)^r + 1}{r} - \sum_{r=1}^{\langle -1/4 \rangle_p} \frac{1}{r} = \begin{cases} H_{[\frac{p}{8}]} - H_{[\frac{p}{4}]} & \text{if } p \equiv 1 \pmod{4}, \\ H_{[\frac{3p}{8}]} - H_{[\frac{3p}{4}]} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

From [S1, Theorems 3.3 and 3.4] we know that

$$\begin{aligned} H_{[\frac{p}{8}]} &\equiv -4q_p(2) - 2 \frac{P_{p-(\frac{2}{p})}}{p} \pmod{p}, \\ H_{[\frac{3p}{8}]} &= H_{[\frac{p}{8}]} + \sum_{\frac{p}{8} < k < \frac{3p}{8}} \frac{1}{k} \equiv H_{[\frac{p}{8}]} + 4 \frac{P_{p-(\frac{2}{p})}}{p} \equiv -4q_p(2) + 2 \frac{P_{p-(\frac{2}{p})}}{p} \pmod{p}. \end{aligned}$$

We also have $H_{[\frac{3p}{4}]} = H_{p-1} - \sum_{k=1}^{[p/4]} \frac{1}{p-k} \equiv H_{[\frac{p}{4}]} \equiv -3q_p(2) \pmod{p}$. Thus,

$$\sum_{r=1}^{\langle -1/4 \rangle_p} \frac{(-1)^r}{r} = \begin{cases} H_{[\frac{p}{8}]} - H_{[\frac{p}{4}]} \equiv -q_p(2) - 2\frac{P_{p-(\frac{2}{p})}}{p} \pmod{p} & \text{if } p \equiv 1 \pmod{4}, \\ H_{[\frac{3p}{8}]} - H_{[\frac{3p}{4}]} \equiv -q_p(2) + 2\frac{P_{p-(\frac{2}{p})}}{p} \pmod{p} & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

Taking $a = -1/4$ in (3.1) we see that $t = ((-1)^{\frac{p-1}{2}} - 2)/4$ and so

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{4k}{2k}}{64^k} \\ &= \sum_{k=0}^{(p-1)/2} \binom{-1/4}{k} \binom{-3/4}{k} \\ &\equiv \begin{cases} (-1)^{\langle a \rangle_p} (1 + 2pt(-2)\frac{P_{p-(\frac{2}{p})}}{p}) = (-1)^{\frac{p-1}{4}} (1 + P_{p-(\frac{2}{p})}) \pmod{p} & \text{if } p \equiv 1 \pmod{4}, \\ (-1)^{\langle a \rangle_p} (1 + 2p(t+1) \cdot 2\frac{P_{p-(\frac{2}{p})}}{p}) = (-1)^{\frac{p-3}{4}} (1 + P_{p-(\frac{2}{p})}) \pmod{p} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \end{aligned}$$

This proves the theorem.

It is well known that ([MOS]) for any positive integer n ,

$$(3.2) \quad \begin{aligned} B_{2n}\left(\frac{1}{2}\right) &= (2^{1-2n} - 1)B_{2n}, \quad B_{2n}\left(\frac{1}{3}\right) = \frac{3 - 3^{2n}}{2 \cdot 3^{2n}} B_{2n}, \\ B_{2n}\left(\frac{1}{4}\right) &= \frac{2 - 2^{2n}}{4^{2n}} B_{2n}, \quad B_{2n}\left(\frac{1}{6}\right) = \frac{(2 - 2^{2n})(3 - 3^{2n})}{2 \cdot 6^{2n}} B_{2n}. \end{aligned}$$

Let $p > 3$ be a prime. From (3.2) and the well-known fact $pB_{p-1} \equiv p-1 \pmod{p}$ (see [IR]) one can easily deduce the following known congruences (see [L],[GS]):

$$(3.3) \quad B_{p-1}\left(\frac{1}{2}\right) - B_{p-1} \equiv 2q_p(2) \pmod{p}, \quad B_{p-1}\left(\frac{1}{3}\right) - B_{p-1} \equiv \frac{3}{2}q_p(3) \pmod{p},$$

$$(3.4) \quad B_{p-1}\left(\frac{1}{4}\right) - B_{p-1} \equiv 3q_p(2) \pmod{p},$$

$$(3.5) \quad B_{p-1}\left(\frac{1}{6}\right) - B_{p-1} \equiv 2q_p(2) + \frac{3}{2}q_p(3) \pmod{p}.$$

In [GS] Graville and Sun showed that

$$(3.6) \quad B_{p-1}\left(\frac{1}{12}\right) - B_{p-1} \equiv 3\frac{S_{p-(\frac{3}{p})}}{p} + 3q_p(2) + \frac{3}{2}q_p(3) \pmod{p},$$

where $\{S_n\}$ is given by $S_0 = 0$, $S_1 = 1$ and $S_{n+1} = 4S_n - S_{n-1}$ ($n \geq 1$).

Theorem 3.3. *Let $p > 3$ be a prime. Then*

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{3k}{k} \binom{6k}{3k}}{432^k} \equiv (-1)^{\frac{p-1}{2}} \left(1 + \left(\frac{3}{p}\right) S_{p-(\frac{3}{p})}\right) \pmod{p^2}.$$

Proof. It is well known that ([MOS])

$$(3.7) \quad E_n(x) = \frac{2}{n+1} \left(B_{n+1}(x) - 2^{n+1} B_{n+1}\left(\frac{x}{2}\right) \right) = \frac{2^{n+1}}{n+1} \left(B_{n+1}\left(\frac{x+1}{2}\right) - B_n\left(\frac{x}{2}\right) \right).$$

For $a \in \mathbb{Z}_p$ we know that $B_{p-1}(a) - B_{p-1} \in \mathbb{Z}_p$ and $pB_{p-1} \equiv p-1 \pmod{p}$ (see [IR,S2,S3]). Thus,

$$(3.8) \quad \begin{aligned} E_{p-2}(-a) &= \frac{2}{p-1} \left(B_{p-1}(-a) - 2^{p-1} B_{p-1}\left(-\frac{a}{2}\right) \right) \\ &= \frac{2}{p-1} \left(B_{p-1}(-a) - B_{p-1} - 2^{p-1} \left(B_{p-1}\left(-\frac{a}{2}\right) - B_{p-1} \right) - (2^{p-1} - 1) B_{p-1} \right) \\ &\equiv -2 \left(B_{p-1}(-a) - B_{p-1} - \left(B_{p-1}\left(-\frac{a}{2}\right) - B_{p-1} \right) + q_p(2) \right) \pmod{p}. \end{aligned}$$

Taking $a = -\frac{1}{6}$ in (3.8) and then applying (3.5) and (3.6) we get

$$(3.9) \quad E_{p-2}\left(\frac{1}{6}\right) \equiv -2 \left(B_{p-1}\left(\frac{1}{6}\right) - B_{p-1} - \left(B_{p-1}\left(\frac{1}{12}\right) - B_{p-1} \right) + q_p(2) \right) \equiv 6 \frac{S_{p-(\frac{3}{p})}}{p} \pmod{p}.$$

Now applying (3.1) we deduce that

$$\sum_{k=0}^{(p-1)/2} \binom{-\frac{1}{6}}{k} \binom{-\frac{5}{6}}{k} \equiv \begin{cases} (-1)^{\frac{p-1}{6}} + \frac{p}{6} E_{p-2}\left(\frac{1}{6}\right) \equiv (-1)^{\frac{p-1}{2}} + S_{p-(\frac{3}{p})} \pmod{p^2} & \text{if } 6 \mid p-1, \\ (-1)^{\frac{5p-1}{6}} - \frac{p}{6} E_{p-2}\left(\frac{1}{6}\right) \equiv (-1)^{\frac{p-1}{2}} - S_{p-(\frac{3}{p})} \pmod{p^2} & \text{if } 6 \nmid p-1. \end{cases}$$

This together with (1.1) yields the result.

Theorem 3.4. *Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$, $a \not\equiv 0 \pmod{p}$ and $t = (a - \langle a \rangle_p)/p$. Then*

$$\begin{aligned} &\sum_{k=0}^{(p-1)/2} k \binom{a}{k} \binom{-1-a}{k} \\ &\equiv \begin{cases} (-1)^{\langle a \rangle_p} a(a+1) - \frac{1}{2} p t (2a(a+1) E_{p-2}(-a) + 2a+1) \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ (-1)^{\langle a \rangle_p} a(a+1) - \frac{1}{2} p (t+1) (2a(a+1) E_{p-2}(-a) + 2a+1) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2}. \end{cases} \end{aligned}$$

Proof. By [S11, Lemma 2.4],

$$\begin{aligned} &\sum_{k=0}^n (k - a(a+1)) \binom{a}{k} \binom{-1-a}{k} \\ &= -a(a+1) \binom{a-1}{n} \binom{-2-a}{n} = -a(a+n+1) \binom{a-1}{n} \binom{-1-a}{n}. \end{aligned}$$

Taking $n = \frac{p-1}{2}$ in the above identity and then applying Lemma 2.1 and (3.1) we deduce that

$$\sum_{k=0}^{(p-1)/2} k \binom{a}{k} \binom{-1-a}{k} - a(a+1) \sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k}$$

$$\begin{aligned}
&= -a \left(a + \frac{p-1}{2} + 1 \right) \binom{a-1}{\frac{p-1}{2}} \binom{-1-a}{\frac{p-1}{2}} \\
&\equiv \begin{cases} -a \left(a + \frac{1}{2} \right) \frac{pt}{\langle a \rangle_p} \equiv -\frac{2a+1}{2} pt \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ -a \left(a + \frac{1}{2} \right) \frac{p(t+1)}{\langle a \rangle_p} \equiv -\frac{2a+1}{2} p(t+1) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2}. \end{cases}
\end{aligned}$$

This together with (3.1) yields the result.

Theorem 3.5. *Let p be a prime greater than 3. Then*

$$\sum_{k=0}^{(p-1)/2} \frac{k \binom{2k}{k} \binom{3k}{k}}{27^k} \equiv -\left(\frac{p}{3}\right) \frac{4-3p+2^{p+2}}{54} \pmod{p^2}.$$

Proof. From (3.3), (3.5) and (3.8) we derive that

$$\begin{aligned}
(3.10) \quad E_{p-2} \left(\frac{1}{3} \right) &\equiv -2 \left(B_{p-1} \left(\frac{1}{3} \right) - B_{p-1} - \left(B_{p-1} \left(\frac{1}{6} \right) - B_{p-1} \right) + q_p(2) \right) \\
&\equiv -2 \left(\frac{3}{2} q_p(3) - 2q_p(2) - \frac{3}{2} q_p(3) + q_p(2) \right) = 2q_p(2) \pmod{p}.
\end{aligned}$$

Now taking $a = -\frac{1}{3}$ in Theorem 3.4 and then applying (1.1) and (3.10) we deduce the result.

4. Congruences for $\sum_{k=0}^{\frac{p-1}{2}} \frac{1}{2k+1} \binom{a}{k} \binom{-1-a}{k}$ and $\sum_{k=0}^{\frac{p-1}{2}} \frac{1}{2k-1} \binom{a}{k} \binom{-1-a}{k} \pmod{p^2}$

Lemma 4.1 ([MS]). *Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$, $a \not\equiv 0 \pmod{p}$ and $t = (a - \langle a \rangle_p)/p$. Then*

$$\sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} \frac{2a+1}{2k+1} \equiv 1 + 2t + 4ptq_p(2) + 2pt \sum_{r=1}^{\langle a \rangle_p} \frac{1}{r} + 2p \sum_{\substack{1 \leq r \leq \langle a \rangle_p \\ r > \frac{p}{2}}} \frac{1}{r} \pmod{p^2}.$$

Theorem 4.1 . *Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$, $a \not\equiv 0 \pmod{p}$ and $t = (a - \langle a \rangle_p)/p$. Then*

$$\begin{aligned}
&\sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} \frac{2a+1}{2k+1} \\
&\equiv \begin{cases} 1 + 2t + 4ptq_p(2) - 2pt(B_{p-1}(-a) - B_{p-1}) \pmod{p^2} & \text{if } \langle a \rangle_p \leq \frac{p-1}{2}, \\ 1 + 2t + 4p(t+1)q_p(2) - 2p(t+1)(B_{p-1}(-a) - B_{p-1}) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p-1}{2}. \end{cases}
\end{aligned}$$

Proof. By [S3, Lemma 3.2],

$$\begin{aligned}
(4.1) \quad \sum_{r=1}^{\langle a \rangle_p} \frac{1}{r} &\equiv \sum_{r=1}^{\langle a \rangle_p} r^{p-2} \equiv (-1)^{p-1} \frac{B_{p-1}(-a) - B_{p-1}}{p-1} + (-a + \langle a \rangle_p) B_{p-2} \\
&\equiv -(B_{p-1}(-a) - B_{p-1}) \pmod{p}.
\end{aligned}$$

Thus, for $\langle a \rangle_p \leq \frac{p-1}{2}$ the result follows from Lemma 4.1.

Now assume $\langle a \rangle_p > \frac{p-1}{2}$. As $H_{\frac{p-1}{2}} \equiv -2q_p(2) \pmod{p}$, we see that

$$\sum_{\substack{1 \leq r \leq \langle a \rangle_p \\ r > \frac{p}{2}}} \frac{1}{r} = \sum_{r=1}^{\langle a \rangle_p} \frac{1}{r} - \sum_{r=1}^{(p-1)/2} \frac{1}{r} \equiv \sum_{r=1}^{\langle a \rangle_p} \frac{1}{r} + 2q_p(2) \pmod{p}.$$

Hence the result follows from the above and Lemma 4.1.

Corollary 4.1. *Let $p > 3$ be a prime. Then*

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{3k}{k}}{(2k+1)27^k} \equiv \left(\frac{p}{3}\right) (2 - 2^{p+1} + 3^p) \pmod{p^2}.$$

Proof. As $t = -\frac{1}{3}$ or $-\frac{2}{3}$ according as $p \equiv 1 \pmod{3}$ or $p \equiv 2 \pmod{3}$, taking $a = -\frac{1}{3}$ in Theorem 4.1 and then applying (3.3) yields

$$\sum_{k=0}^{(p-1)/2} \frac{1}{2k+1} \binom{-\frac{1}{3}}{k} \binom{-\frac{2}{3}}{k} \equiv \left(\frac{p}{3}\right) (1 - 4pq_p(2) + 3pq_p(3)) = \left(\frac{p}{3}\right) (2 - 2^{p+1} + 3^p) \pmod{p}.$$

This together with (1.1) gives the result.

Theorem 4.2 . *Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$ and $a \not\equiv 0 \pmod{p}$. Then*

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} \frac{1}{2k-1} \\ & \equiv \begin{cases} -(2a+1) + 2(a - \langle a \rangle_p)(1 + (2a+1)E_{p-2}(-2a)) \pmod{p^2} & \text{if } \langle a \rangle_p < \frac{p}{2}, \\ 2a+1 + 2(p+a - \langle a \rangle_p)(1 + (2a+1)E_{p-2}(-2a)) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p}{2}. \end{cases} \end{aligned}$$

Proof. By [S10, Lemma 3.1],

$$\begin{aligned} (4.2) \quad & \sum_{k=0}^n \binom{a}{k} \binom{-1-a}{k} \frac{(2a(a+1)+1)k - a(a+1)}{4k^2-1} \\ & = \frac{a(a+1)}{2n+1} \binom{a-1}{n} \binom{-2-a}{n} = \frac{a(a+n+1)}{2n+1} \binom{a-1}{n} \binom{-a-1}{n}. \end{aligned}$$

Set $t = (a - \langle a \rangle_p)/p$. By Lemma 2.1,

$$\begin{aligned} & \frac{1}{p} \binom{a-1}{\frac{p-1}{2}} \binom{-a-1}{\frac{p-1}{2}} \\ & \equiv \begin{cases} \frac{t}{\langle a \rangle_p} + \frac{pt}{a} (2q_p(2) + H_{\frac{p-1}{2} - \langle a \rangle_p}) - \frac{pt^2}{a^2} \pmod{p^2} & \text{if } \langle a \rangle_p < \frac{p}{2}, \\ \frac{t+1}{\langle a \rangle_p} + \frac{p(t+1)}{a} (2q_p(2) + H_{\langle a \rangle_p - \frac{p+1}{2}}) - \frac{pt(t+1)}{a^2} \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p}{2}. \end{cases} \end{aligned}$$

For $\langle a \rangle_p < \frac{p}{2}$, using (4.1) and the fact $B_n(1-x) = (-1)^n B_n(x)$ ([MOS]) we see that

$$H_{\frac{p-1}{2}-\langle a \rangle_p} \equiv -\left(B_{p-1}\left(\frac{1}{2}+a\right)-B_{p-1}\right) = -\left(B_{p-1}\left(\frac{1}{2}-a\right)-B_{p-1}\right) \pmod{p}.$$

For $\langle a \rangle_p > \frac{p}{2}$, using (4.1) we see that

$$H_{\langle a \rangle_p - \frac{p+1}{2}} \equiv -\left(B_{p-1}\left(\frac{1}{2}-a\right)-B_{p-1}\right) \pmod{p}.$$

Thus, taking $n = \frac{p-1}{2}$ in (4.2) and applying the above we obtain

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} \frac{(2a(a+1)+1)k - a(a+1)}{4k^2-1} \\ & \equiv \begin{cases} \left(a + \frac{p+1}{2}\right) \left(\frac{t(pt + \langle a \rangle_p)}{\langle a \rangle_p} + pt(2q_p(2) + B_{p-1} - B_{p-1}(\frac{1}{2}-a)) - \frac{pt^2}{a}\right) \pmod{p^2} \\ \quad \text{if } \langle a \rangle_p < \frac{p}{2}, \\ \left(a + \frac{p+1}{2}\right) \left(\frac{(t+1)(pt + \langle a \rangle_p)}{\langle a \rangle_p} + p(t+1)(2q_p(2) + B_{p-1} - B_{p-1}(\frac{1}{2}-a))\right. \\ \quad \left. - \frac{pt(t+1)}{a}\right) \pmod{p^2} \quad \text{if } \langle a \rangle_p > \frac{p}{2} \end{cases} \\ & \equiv t' \left(\frac{p}{2} + \frac{2a+1}{2} \left(1 + p(2q_p(2) + B_{p-1} - B_{p-1}(\frac{1}{2}-a))\right)\right) \pmod{p^2}, \end{aligned}$$

where $t' = t$ or $t+1$ according as $\langle a \rangle_p < \frac{p}{2}$ or $\langle a \rangle_p > \frac{p}{2}$. As

$$\frac{1}{2k-1} = 4 \frac{(2a(a+1)+1)k - a(a+1)}{4k^2-1} - (2a+1) \frac{2a+1}{2k+1},$$

from the above and Theorem 4.1 we see that

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \frac{1}{2k-1} \binom{a}{k} \binom{-1-a}{k} \\ & = 4 \sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} \frac{(2a(a+1)+1)k - a(a+1)}{4k^2-1} \\ & \quad - (2a+1) \sum_{k=0}^{(p-1)/2} \binom{a}{k} \binom{-1-a}{k} \frac{2a+1}{2k+1} \\ & \equiv 2pt' + 2(2a+1)t' \left(1 + p(2q_p(2) + B_{p-1} - B_{p-1}(\frac{1}{2}-a))\right) \\ & \quad - (-1)^{t'-t} (2a+1) - 2(2a+1)t' (1 + p(2q_p(2) - B_{p-1}(-a) + B_{p-1})) \\ & = -(-1)^{t'-t} (2a+1) + 2pt' + 2(2a+1)pt' \left(B_{p-1}(-a) - B_{p-1}(\frac{1}{2}-a)\right) \pmod{p^2}. \end{aligned}$$

By (3.7),

$$(4.3) \quad E_{p-2}(-2a) = \frac{2^{p-1}}{p-1} \left(B_{p-1}\left(-a + \frac{1}{2}\right) - B_{p-1}(-a) \right) \equiv B_{p-1}(-a) - B_{p-1}\left(\frac{1}{2} - a\right) \pmod{p}.$$

Now combining the above gives the result.

Theorem 4.3. *Let p be a prime greater than 3. Then*

$$\begin{aligned} \text{(i)} \quad & \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{4k}{2k}}{(2k-1)64^k} \equiv (-1)^{\frac{p+1}{2}} \frac{p+1}{2} \pmod{p^2}, \\ \text{(ii)} \quad & \sum_{k=0}^{(p-1)/2} \frac{\binom{2k}{k} \binom{3k}{k}}{(2k-1)27^k} \equiv \frac{1}{9} \left(\frac{p}{3}\right) (2^{p+1} - 7 - 6p) \pmod{p^2}, \\ \text{(iii)} \quad & \sum_{k=0}^{(p-1)/2} \frac{\binom{3k}{k} \binom{6k}{3k}}{(2k-1)432^k} \equiv -\frac{1}{9} \left(\frac{p}{3}\right) (2^{p+1} + 2 + 3p) \pmod{p^2}. \end{aligned}$$

Proof. It is well known that ([MOS])

$$B_n(1-x) = (-1)^n B_n(x) \quad \text{and} \quad E_n(1-x) = (-1)^n E_n(x).$$

For $m = 3, 4, 6$ we see that $\langle -\frac{1}{m} \rangle_p = \frac{p-1}{m}$ or $\frac{(m-1)p-1}{m}$ according as $p \equiv 1 \pmod{m}$ or not. Using (4.3) we see that $E_{p-2}(\frac{1}{2}) \equiv B_{p-1}(\frac{1}{4}) - B_{p-1}(\frac{3}{4}) \equiv 0 \pmod{p}$. Now taking $a = -\frac{1}{4}$ in Theorem 4.2 and then applying (1.1) yields (i). Using (3.10) we see that $E_{p-2}(\frac{2}{3}) = -E_{p-2}(\frac{1}{3}) \equiv -2q_p(2) \pmod{p}$. Taking $a = -\frac{1}{3}$ in Theorem 4.2 and then applying (1.1) yields (ii). Taking $a = -\frac{1}{6}$ in Theorem 4.2 and then applying (3.10) and (1.1) yields (iii). The proof is now complete.

Theorem 4.4. *Let $p > 3$ be a prime, $a \in \mathbb{Z}_p$, $a \not\equiv 0 \pmod{p}$ and $t = (a - \langle a \rangle_p)/p$. Then*

$$\sum_{k=0}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{2k-1} \equiv \begin{cases} 2pt - (2a+1) \sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \pmod{p^2} & \text{if } \langle a \rangle_p < \frac{p}{2}, \\ 2p(t+1) - \frac{2a+1}{2t+1} - (2a+1) \left(1 + \frac{1}{2t+1}\right) \sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p}{2} \text{ and } t \not\equiv -\frac{1}{2} \pmod{p}, \\ 2a+1 + p(1 + (2a+1)E_{p-2}(-2a)) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p}{2} \text{ and } t \equiv -\frac{1}{2} \pmod{p}. \end{cases}$$

Proof. By (1.9),

$$\sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \equiv (-1)^{\langle 2a \rangle_p} - (2a - \langle 2a \rangle_p) E_{p-2}(-2a)$$

$$= \begin{cases} 1 - (2a - 2\langle a \rangle_p) E_{p-2}(-2a) \pmod{p^2} & \text{if } \langle a \rangle_p < \frac{p}{2}, \\ -1 - (2a - 2\langle a \rangle_p + p) E_{p-2}(-2a) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p}{2}. \end{cases}$$

Thus, from Theorem 4.2 we deduce that

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{2k-1} + (2a+1) \sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \\ & \equiv \begin{cases} 2(a - \langle a \rangle_p) \pmod{p^2} & \text{if } \langle a \rangle_p < \frac{p}{2}, \\ 2(p + a - \langle a \rangle_p) + (2a+1)p E_{p-2}(-2a) \pmod{p^2} & \text{if } \langle a \rangle_p > \frac{p}{2}. \end{cases} \end{aligned}$$

This shows that the result is true for $\langle a \rangle_p < \frac{p}{2}$. Now assume $\langle a \rangle_p > \frac{p}{2}$. From the above we have

$$1 + \sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \equiv -(2t+1)p E_{p-2}(-2a) \pmod{p^2}.$$

Hence for $t \not\equiv -\frac{1}{2} \pmod{p}$,

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{2k-1} + (2a+1) \sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \\ & \equiv 2(p + a - \langle a \rangle_p) - \frac{2a+1}{2t+1} \left(1 + \sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \right) \pmod{p^2}. \end{aligned}$$

This yields the result in this case. For $t \equiv -\frac{1}{2} \pmod{p}$, from the above we have $a - \langle a \rangle_p = pt \equiv -\frac{p}{2} \pmod{p^2}$ and $\sum_{k=0}^{p-1} \binom{2a}{k} (-2)^k \equiv -1 \pmod{p^2}$. Hence

$$\begin{aligned} & \sum_{k=0}^{(p-1)/2} \frac{\binom{a}{k} \binom{-1-a}{k}}{2k-1} \equiv 2a+1 + 2(p + a - \langle a \rangle_p) + (2a+1)p E_{p-2}(-2a) \\ & \equiv 2a+1 + p(1 + (2a+1) E_{p-2}(-2a)) \pmod{p^2}. \end{aligned}$$

This yields the result in this case. The proof is now complete.

Acknowledgment

The author is supported by the National Natural Science Foundation of China (grant No. 11371163).

References

- [EMOT] A. Erdélyi, W. Magnus, F. Oberhettinger and F.G. Tricomi, *Higher Transcendental Functions, Vol.I*, McGraw-Hill, New York, 1953. Based, in part, on notes left by Harry Bateman.

- [GS] A. Granville and Z.W. Sun, *Values of Bernoulli polynomials*, Pacific J. Math. **172**(1996), 117-137.
- [IR] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, second ed., Springer, New York, 1990.
- [L] E. Lehmer, *On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson*, Ann. of Math. **39**(1938) 350-360.
- [MOS] W. Magnus, F. Oberhettinger and R.P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics (3rd edn.)* (Springer, New York, 1966), pp. 25-32.
- [MS] G.S. Mao and Z.W. Sun, *New congruences involving products of two binomial coefficients*, arXiv:1601.04782, 2016.
- [M1] E. Mortenson, *A supercongruence conjecture of Rodriguez-Villegas for a certain truncated hypergeometric function*, J. Number Theory **99**(2003), 139-147.
- [M2] E. Mortenson, *Supercongruences between truncated ${}_2F_1$ hypergeometric functions and their Gaussian analogs*, Trans. Amer. Math. Soc. **355**(2003), 987-1007.
- [RV] F. Rodriguez-Villegas, *Hypergeometric families of Calabi-Yau manifolds* in: Noriko Yui, James D. Lewis (Eds.), *Calabi-Yau Varieties and Mirror Symmetry*, Toronto, ON, 2001, in: Fields Inst. Commun., vol. 38, Amer. Math. Soc., Providence, RI, 2003, pp.223-231.
- [S1] Z.H. Sun, *The combinatorial sum $\sum_{k \equiv r \pmod{m}} \binom{n}{k}$ and its application in number theory III*, J. Nanjing Univ. Math. Biquarterly **12**(1995), 90-102.
- [S2] Z.H. Sun, *Congruences for Bernoulli numbers and Bernoulli polynomials*, Discrete Math. **163**(1997) 153-163.
- [S3] Z.H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105**(2000) 193-223.
- [S4] Z.H. Sun, *Congruences involving Bernoulli polynomials*, Discrete Math. **308**(2008) 71-112.
- [S5] Z.H. Sun, *Congruences involving Bernoulli and Euler numbers*, J. Number Theory **128**(2008) 280-312.
- [S6] Z. H. Sun, *Identities and congruences for a new sequence*, Int. J. Number Theory **8**(2012) 207-225.
- [S7] Z. H. Sun, *Some properties of a sequence analogous to Euler numbers*, Bull. Aust. Math. Soc. **87**(2013) 425-440.
- [S8] Z.H. Sun, *Generalized Legendre polynomials and related supercongruences*, J. Number Theory **143**(2014) 293-319.

- [S9] Z.H. Sun, *Super congruences concerning Bernoulli polynomials*, Int. J. Number Theory **11**(2015), 2393-2404.
- [S10] Z.H. Sun, *Supercongruences involving Bernoulli polynomials*, Int. J. Number Theory **12**(2016), 1259-1271.
- [S11] Z.H. Sun, *Supercongruences involving Euler polynomials*, Proc. Amer. Math. Soc. **144**(2016), 3295-3308.
- [Su1] Z.W. Sun, *Super congruences and Euler numbers*, Sci. China Math. **54**(2011), 2509-2535.
- [Su2] Z.W. Sun, *p-adic congruences motivated by series*, J. Number Theory **134**(2014) 181-196.
- [T1] R. Tauraso, *Congruences involving alternating multiple harmonic sums*, Electron. J. Combin. **17**(2010), #R16, 11 pp.
- [T2] R. Tauraso, *Supercongruences for a truncated hypergeometric series*, Integers **12**(2012), #A45, 12 pp.